

< 0.25 mm mesh and were oscillated vertically in water for twenty (20) minutes at the rate of one oscillation per second. After wet-sieving, the remaining soil materials on each sieve were emptied into beakers, dried for 48 hours in an oven and weighed (W2). The percentage of sand in each sieve fractions was corrected by washing the aggregates with sodium hexametaphosphate, oven dried and weighed (W3). Percentage water stable aggregates (%WSA) was then calculated using $\% \text{WSA} = \frac{W2-W3}{W1-W3} \times 100$Equ. 1

The same procedure apart from oven drying the soil in each sieve was used to obtain soil samples for chemical analyses in soil aggregate sizes.

Bouyoucos hydrometer method documented by Gee and Or, (2002) was used in the determination of particle size. Core method of Grossman and Reinsch, (2002) was used in bulk density determination. The method documented by Hendershot et al., (1993) was used in the measurement of soil pH in soil-water ratio of 1:2.5. Soil organic carbon in whole soil and in subsamples of aggregate sizes was measured by wet digestion method of Nelson and Sommers, (1996). Available phosphorus was measured by Olsen method (Emteryd, 1989).

The use of flame photometer according to Thomas (1982) aided exchangeable cations (potassium and sodium) determination while magnesium and calcium were determined using ethylene diamine tetra-acetic acid (EDTA) (Thomas, 1982). Effective cation exchange capacity was determined by summing the exchangeable cations as documented by Landon (1991) Brady and Weil (1999).

Data Analyses. Data were analysed with analysis of variance ANOVA. Significant means were separated using least significant difference (LSD) at 5% probability.

Results and Discussion

The results of the physical and chemical properties of the studied soils are presented in Table 1. Sand fraction ranged from 552.90 to 840.00 gkg^{-1} and is significantly higher ($p < 0.05$) in Uzoakoli. Clay fraction ranged from 110.00 gkg^{-1} to 323.30 gkg^{-1} , while silt ranged from 40.00 gkg^{-1} to 123.80 gkg^{-1} . These values were typical of soils of south eastern Nigeria (Igwe, 1995). Values of silt fraction which followed the order: ARG > FBS > CPS > BAG are low indicating high degree of weathering. Texturally, soil of Ishiagu was sandy clay loam, Obinze: sandy, Umuna and Uzoakoli were sandy loam and loamy sand respectively. The classification was in line with Esheth (1995), Akamigbo and Asadu (1983) who documented similar textures for surface horizon soils of humid tropics. Sandiness of texture could be attributed to land use, climate and geology. Soils with high sand fraction witness leaching, of high infiltration and low water retention thus incapable of supporting most crops. Differences in clay fraction are attributed to soil structure, clay movement and parent material. Bulk density ranged from 1.20 gcm^{-3} in Uzuakoli to 1.40 gcm^{-3} in Obinze. Soils whose bulk densities were low are related to high porosity, lesser compaction and easier penetration of roots and movement of micro fauna in them. Higher bulk density of soils leads to low infiltration and high runoff. However these values were lower than 1.85 gcm^{-3} (SSS 2006) considered as very critical and will surely impair root growth and is tandem with those of tropical soils (Landon, 1991).

The stability of aggregates using percentage water stable aggregates greater 0.25 mm (WSA > 0.25) was significantly higher in soil of Umuna under FBS (59.18 %) and least at

soils of Obinze (26.27 %) under Coastal Plain Sand. The trend in the stability with respect to studied parent material was FBS >ARG>BAG> CPS (Table 1). This trend is attributed to texture, type of clay (USDA 1996) and organic matter which has binding effects soil particles (Tisdall and Oades, 1982; Hayness and Swift, 1990; USDA, 1996) and their interactions (Le-Bissonnaise, 1996). Others include exchangeable sodium percentage (Kazmanet *al.*, 1983), soil management (Iraj, 2009). Kay and Angers (1999) stated that aggregate stability declines rapidly as organic carbon decreases from 1.5- 1.2%. Among the list of other factors that affect structural stability include sesquioxides (Le-Bissonnaise and Singer 1993). This result showed that soil under False bedded sand stone is much more stable than others and will be able to resist erosion more than others. Soil pH in water though did not differ significantly ($p < 0.05$) among parent materials ranged from 5.37 to 5.75 (Table 1). Moderate acidic condition of the soils was credited to constant leaching by rainfall in the area. This leaves the soil saturated with more aluminium and hydrogen (Landon, 1991).

The result of soil organic carbon presented in Table 1 significantly differed ($p < 0.05$) with parent material and followed similar trend with that of aggregate stability (FBS >ARG>BAG> CPS). Soil organic carbon (SOC) ranged from 9.60 in Obinze - 14.80 gkg^{-1} in soils under Uzuakoli. Generally, the result of soil organic matter are rated low (Landon 1991; Igwe and Stahr 2004, Opara - Nadi (1988), typical of tropical soils, a factor that predisposes the soil to dispersion making it prone to erosion (Agim et al., 2012).

Table 1. Physical properties of studied soil.

P.M	Location	Sand	Silt gkg^{-1}	Clay	T.C	B.D gcm^{-3}	WSA %	pH (H_2O)	SOC gkg^{-1}	Av, P mgkg^{-1}	T.N gkg^{-1}	ECEC cmolkg^{-1}
ARG	Ishiagu	552.90	123.80	323.30	SCL	1.25	47.16	5.37	12.40	10.94	1.20	7.41
CPS	Obinze	809.60	80.40	110.00	S	1.40	26.27	5.75	9.60	11.26	1.02	3.97
FBS	Umuna	622.90	90.40	286.70	SL	1.30	59.18	5.75	10.90	11.57	1.30	12.00
BAG	Uzoakoli	840.00	40.00	120.00	LS	1.20	31.15	5.42	14.80	11.86	1.20	3.21
LSD ($P < 0.05$)		22.10*	7.31*	10.42*		0.21*	2.23*	NS	0.39**	NS	NS	0.95*

P.M=parent material, ARG=Asu River Group, CPS=Coastal plain sand, FBS=False bedded sand, BAG=Bende Ameki Group, WSA=water stable aggregates. BD=Bulk density. *=Significant, **=highly significant, ECEC=effective cation exchange capacity, SOC=Soil organic carbon=probability level, Av. P=Available phosphorus, T.C=Textural class.

The distribution of selected soil nutrients in aggregate sizes of studied soil

The distribution of selected soil properties in aggregate sizes of studied soils are presented in Table 2. Soil organic carbon ranged from 10.31 gkg^{-1} -12.20 gkg^{-1} in aggregate diameters of 2 and 0.5 mm in Ishiagu (ARG), 5.30 - 9.30 gkg^{-1} in 2 and < 0.25 mm in Obinze (CPS). Similarly, values of 8.80 - 10.60 and 8.40 - 14.30 gkg^{-1} were obtained in Umuna (FBS) and Uzoakoli (BAG) under sieve sizes of 2, < 0.25 and 0.5 - < 0.25 mm respectively. Generally, these values were lower than that contained in whole soil (Figures 1:a-d). This confirms that breakdown of aggregates leads to loss of organic matter. In all cases however, organic matter were low, highest values were obtained in the 0.5 mm sieve at Ishiagu, where as they occurred at the <0.25 mm for others. Result obtained in Ishiagu is similar to that of Igwe 2001, Green et al., (2005) where as those of others followed the trend of Razalfimbelo et

al., (2008) who reported high SOC in micro aggregates (<0.02 mm). Inconsistent values are attributed to parent materials. The high values of soil organic matter at this level of aggregation are also attributed to slaking which occur when bigger aggregates are not able to survive inner pressure as a result of fast water absorption or tillage. Such process releases heat that makes the aggregate to collapse. Igwe (2003) observed that soils whose aggregates occur more on the < 0.25 mm sizes correlate with inter rill and rill erosion in the field. Le Bissoin (1996) posited that aggregates at these sizes are very unstable and undesirable. This implies that the organic matter contained in the above soils are very much prone to erosion losses. This also supports Mbagwu et al., (1993), Obi (1982), who found significant positive association between micro aggregate stability and organic matter concentration in tropical soils.

Total nitrogen followed the same order with organic carbon. Higher values were noted in 0.5 and in < 0.25 mm sieve sizes. Wright and Inglett (2009) found similar trends in organic nitrogen distribution in aggregates of Everglades Histosols. Guan et al., (2018) also found similar result in an alpine meadow at the Damxung Grassland Tibetan Autonomous Region, northwest China. Increase in values of nitrogen at this level of aggregation is an indication of high preservability of soil nutrient micro aggregates or clay sized fractions.

Available phosphorus was significantly ($p < 0.05$) higher in aggregates size of 1mm in all studied soils except in Uzuakoli. It ranged from 2.90 – 5.70 mg kg^{-1} in sieve size of 2mm and 1mm in Ishiagu, 3.84 – 6.64 mg kg^{-1} in 0.5 and 2 mm sieves in Obinze, 6.60 – 9.60 mg kg^{-1} in sieve size of 0.25 and 2mm in Umuna and 0.10 to 9.40 mg kg^{-1} in aggregates sizes of 0.5 and 0.25mm in Uzoakoli respectively. Available phosphorus decreased as sieve sizes tends to < 0.25mm though not in all cases (Figures 2a and b). Aguilar and Heil (1988) observed that variations in phosphorus content along a toposequence reflected differences in parent material or geology. These values were low (Landon, 1991, Enwezor 1977) and typical of soils of the eastern Nigeria. Uzoho and Oti (2005) recorded similar trend in similar soil. Igwe (2001) attributed low phosphorus availability in soils of the area to high fixation associated with the studied soils. Low phosphorus availability in tropical soil is also attributed to intense weathering and partly to the low availability of phosphorus in the aluminium and iron combination which were the dominant sources in these soils (Brady and Weil 1999). It has also been documented that low phosphorus values inhibit effective nodulation and leads to biological nitrogen fixation process of leguminous plants (Brady and Weil 1999).

Effective cation exchange capacity (ECEC) ranged from 5.77 to 9.03 cmol kg^{-1} in sieves of 2mm and 1mm in Ishiagu (ARG), from 2.20 to 2.93 cmol kg^{-1} in sieves of < 0.25 mm and 1mm, Obinze (CPS); from 5.22 to 11.13 cmol kg^{-1} in sieves of 2mm and 0.5 mm in Umuna (FBS) and from 3.04 to 3.44 cmol kg^{-1} in sieves of < 0.25 and 2 mm in Uzoakoli (BAG) respectively Table 2. Trends in effective cation exchange capacity differed significantly among parent materials suggesting that the breakdown of macro aggregates to smaller fractions during cultivation may lead to significant loss of cations. Generally higher values were obtained macro aggregates 2-0.5 mm than in micro aggregates <0.25. This accounts for higher stability of aggregates at macro levels a fit attributed to the binding actions of cations. Moreover, cations especially divalent ones are tightly held at the exchange complexes of macro aggregates (Jiang et al., 2011) found higher calcium and magnesium at the macro levels of stability in their work. They however obtained high levels of sodium at <0.25 mm

confirming high level of dispersibility thus favouring erosion in the soils. ECEC like soil organic carbon in all soils are low. These values are below the critical limits for soils of South-eastern Nigeria (Enwezor et al., 1990), suggesting poor fertility status of the soil (Igwe, 2000). Ojanuga and Awojuola (1981) attributed low ECEC to the type of clay minerals prominent in the study area and differences in parent materials.

Table 2. Distribution of selected properties in different aggregate sizes

Parent material	Location	Sieve Sizes mm	pH (H ₂ O)	SOC gkg ⁻¹	Av.P mgkg ⁻¹	Total nitrogen gkg ⁻¹	ECEC cmolkg ⁻¹
Asu River Group	Ishiagu	2.00		10.31		0.99	
			5.49		2.90		5.77
		1.00	5.67	11.39	5.70	0.98	9.03
		0.50	5.56	12.20	5.20	1.11	7.37
		<0.25	5.51	11.00	4.50	1.18	5.79
LSD(P< 0.05)		NS	0.08*	0.21*	0.02*	1.18*	
PM X Ss		NS	0.16*	0.05*	0.01*	NS	
Coastal plain sand	Obinze	2.00	5.63	5.30	6.64	0.98	2.55
		1.00	5.83	7.30	4.06	1.01	2.93
		0.50	5.58	8.30	3.84	1.02	2.43
		<0.25	5.85	9.30	4.01	1.02	2.20
		LSD (P <0.05)		0.23*	0.12*	1.00*	NS
PM X Ss		0.05*	0.32*	NS	NS	1.22*	
Falsebedded sand	Umuna	2.00		8.80		1.20	
			5.41		8.60		5.22
		1.00	5.26	9.10	9.60	1.10	8.59
		0.50	5.32	10.50	9.00	1.28	11.13
		<0.25	5.48	10.60	6.60	1.29	9.18
LSD (P < 0.05)		0.27*	NS	0.03*	NS	2.14*	
PM X Ss		1.23*	1.23*	1.10*	NS	2.16*	
Bende Ameki Group	Uzuakoli	2.00		14.00		1.15	
			5.54		0.20		3.44
		1.00	5.71	9.80	3.90	1.10	3.15
		0.50	5.09	8.40	0.10	1.10	3.16
		<0.25	5.04	14.30	9.40	0.09	3.04
LSD (P <0.05)		1.32*	0.17*	0.31*	NS	NS	
PM X Ss		1.20*	1.76*	1.77*	NS	NS	

ARG=Asu River Group,CPS=Coastal plain sand, FBS=Falsebedded sand,BAG=Bende Ameki Group,SOC= Soil organic carbon, ECEC =Effective cation exchange capacity *=Significant,**=highly significant, P= probability level, LSD=Least significant difference. PM x Ss= Interaction between parent material and sieve sizes, Av.P=Available phosphorus




Figure 1a. Soil organic carbon in whole soil before sieving and in sieve sizes under the Asu River Group formation.




Figure 1b. Soil organic carbon in whole soil before sieving and in sieve sizes under Coastal Plain Sand.




Figure 1c. Soil organic carbon in whole soil before sieving and in sieve sizes under the Falsebedded Sand Stone




Figure 1d. Soil organic carbon in whole soil before sieving and in sieve sizes under Bende Ameki Group.




Figure 2a. Available phosphorus in whole soil before sieving and in sieve sizes under Asu River Group and Coastal Plain Sand.




Figure 2b. Available phosphorus in whole soil before sieving and in sieve sizes under Falsbedded sand and Bende Ameki Group

Relationships among selected soil properties of studied soil.

In the result, soil organic carbon had significant ($P < 0.05$) positive relation with water stable aggregates WSA ($r = 0.50$), pH water ($r = 0.53$), total nitrogen ($r = 0.79$), effective cation exchange capacity ($r = 0.51$) and negatively related with bulk density ($r = -0.58$) Table 3. This implies that increase in organic matter content of the soil increases soil stability, soil pH, total nitrogen and effective cation exchange capacity by 50, 53 and 79 and 51 percents respectively. These results were similar to Agim et al., (2012) Sakin (2012) and confirm that of Unamba – Opara, (1982) on selected Nigerian soils. Increase in the structural stability of

the soil was as the result of binding effects of organic matter on soil properties. While organic matter gives flesh to the sand content remain the skeleton (Brady and Weil, 1999). The content of organic matter and nitrogen in soils are interwoven hence the positive relationship among them (Sakin, 2012; Unamba-Opara, 1982). This is because microbial decomposition of organic matter leads to nitrogen availability in soils. This assertion is not ignorant of the fact that nitrogen availability in soil is also achieved via synthetic fertilizer application and atmospheric fixation. Result of the relationship between organic carbon and effective cation exchange capacity supports the fact that it is a reservoir of soil cations and therefore availability of micronutrients in the soil are imminent. Onwudike et al., (2016) attributed positive relations between soil organic matter, micronutrients and soil pH to chelating characteristics of organic matter which helps to hold nutrients in the soil. In his own Verma et al., (2005) attributed such positive correlation to complexing agents that are generated by organic matter which promotes nutrient availability in soil. Negative relationship between soil organic matter and bulk density confirms is in order. This is because microbial decomposition of organic matter increases soil pores, thus low bulk density.

Total nitrogen related positively with moisture content and aggregate stability ($r = 0.38$, and 0.50 , $P < 0.05$). This goes further to buttress the important of organic matter in soil.

Table 3. Correlation matrix among selected properties of studied soil.

	Units	Sand	Silt	Clay	MC	BD	WSA	PH	SOC	TN	Av.p	ECEC
Sand	gkg ⁻¹	*										
Silt	gkg ⁻¹	0.29 ^{ns}	*									
Clay	gkg ⁻¹	0.98**	0.15 ^{ns}	*								
MC	gkg ⁻¹	0.48*	0.01 ^{ns}	0.57*	*							
BD	gcm ⁻³	0.46*	0.01 ^{ns}	0.52*	0.53*	*						
WSA	%	0.53*	0.05 ^{ns}	0.59*	0.89*	0.74*	*					
pH		-0.29 ^{ns}	0.11 ^{ns}	0.29 ^{ns}	0.01 ^{ns}	0.06 ^{ns}	0.16 ^{ns}	*				
SOC	%	0.22 ^{ns}	0.06 ^{ns}	0.30*	0.76*	-0.58**	0.73 ^{ns}	0.53 ^{ns}	*			
TN	%	0.15 ^{ns}	0.06 ^{ns}	0.20 ^{ns}	0.38*	0.10 ^{ns}	0.50*	0.03 ^{ns}	0.79*	*		
Av.P	mgkg ⁻¹	0.07 ^{ns}	0.07 ^{ns}	0.07 ^{ns}	0.03 ^{ns}	0.11 ^{ns}	0.34*	0.04 ^{ns}	0.02 ^{ns}	0.02 ^{ns}	*	
ECEC	cmolk ⁻¹	0.66*	0.37 ^{ns}	0.60*	0.11 ^{ns}	0.28 ^{ns}	0.08 ^{ns}	0.31 ^{ns}	0.51*	0.09 ^{ns}	0.08 ^{ns}	1

Conclusion

In conclusion results showed that variations occur in soil nutrients among geologic formations and aggregates. Organic carbon, available phosphorus and effective cation exchange capacity of soil varied with sieve sizes. Soil organic carbon was lesser in the aggregate sizes in comparison to that of the whole soil at the beginning of the study. The stability of aggregates using percentage water stable aggregates greater 0.25 mm (WSA > 0.25) was followed the trend: FBS>ARG>BAG>CPS. Effective cation exchange capacity followed the trend ARG>FBS>CPS>BAG. Significant positive relations were noted between soil organic carbon and selected properties. Soil organic carbon is very important in the protection of aggregates from deformation, therefore efforts including cover cropping,

mulching application of organic amendment etc that improve organic matter content of the soil should be adopted. Again since soil erosion is a function of rainfall, measures such as afforestation, mulching etc that will dissipate the kinetic energy of falling rain drops before it strikes the soil surface and those that would be able to trap sediments on site are recommended.

Acknowledgments

The work is self-funded. The authors appreciate the Institute of erosion studies, staff of the Department of Soil Science Federal Government certified laboratory all in Federal University of Technology Owerri and Mrs Chinenye Agim for their efforts toward the successful completion of this work.

References

- Agim, L. C., 2016. Structural stability, erodibility and organic amendment effects on soils of some geologic formations in South eastern Nigeria. PhD thesis submitted to Faculty of Agriculture, University of Nigeria Nsukka, 1-277.
- Agim L.C., G.E. Osuji., E.U. Onweremadu., B.N. Ndukwu and S.K. Osuaku, 2012. Seasonal dynamics of soil organic matter and total nitrogen in soils under different land uses in Owerri South eastern Nigeria. *Agro Science Journal of tropical Agriculture, food, environment and Extention*, Vol.1, No.1, 43-54.
- Aguilar, R. and R. D. Heil, 1988. Soil organic carbon, nitrogen and phosphorus quantities in Northern Great Plains Rangeland. *Soil Sci. Soc. Am. J.*, Vol. 52, 1076-1081.
- Akamigbo F.O.R., C.L.A. Asadu, 1983. Influence of parent material on the soil southeastern Nigeria. *East Africa Agricultural Forestry, Journal*, Vol. 48, 82-91.
- Anyadike, R.N.C., 2002. Climate and Vegetation p. 73-8. In G.E.K. Ofomata (ed.). *A survey of Igbo Nation*. Africana Publishers Ltd, Onitsha.
- Brady, N.C and R.R. Weil, 1999. The nature and properties of soil. Twelfth edition. Prince Hall. Upper saddle River, New Jersey, 130 - 265.
- Cambardella, C.A. and E.T. Elliot, 1993. Carbon and nitrogen distribution in Aggregates from cultivated and native grassland soils. *Soil Sci. Soc. Am. J.*, Vol.,57,2071 – 1076.
- Chikezie, I. A., H., Eswaran, D.O., Asawalam, and A.O., Ano, 2010. Characterization of two benchmark soils of contrasting parent materials in Abia State, Southeastern Nigeria. *Global Journal of Pure and Applied Sciences*, Vol.16, No.1., 23-29.
- Cruvinel, P.E., S. Flocchini, J.R. Crestana, J. Morales, B.H. Kusko, and D.R Nielsen, 1990. Studying some influence of the Aggregate Sizes on some Elements of an Oxisol with PIXE. *Soil Sci.*, Vol. 155, 100 – 104.
- De-Swardt A.M, O. P. Casey, 1963. The Coal resources of Nigeria. *Bull. Geol. Survey of Nigeria*: 28. Dixon, J.C, 2015. Soil Morphology in the Critical Zone: The Role of Climate, Geology, and Vegetation in Soil Formation in the Critical Zone. In *Developments in Earth Surface Processes* Volume 19, 147-172. <https://doi.org/10.1016/B978-0-444-63369-9.00005-7>.

Emteryd O., 1989. Chemical and physical analysis of inorganic nutrients in plants, soils, water and air Serial No. Uppsala, Swedish University Agricultural Science.

Enwezor, W.O., 1977. Predicting responses of phosphate applications for soil of south eastern Nigeria. *Soil Sci*, 111-116.

Enwezor, W.O, A.C. Ohiri, E.E. Opoaribo, and E.J. Udo, 1990. A review of soil fertilizer use for crops in southeastern zone of Nigeria (in five volumes). Produced by the Federal Ministry of Agriculture and Natural Resources Lagos.

Esheth E.T., 1995. Chemistry, clay activity and mineralogy of soil along 3 Toposequence over igneous and sedimentary lithology in south eastern Nigeria. *Thai J. Agric Science Vol.2*, 27 - 285.

Eswaran, H., R. Almaraz, E. Van Den Berg and P.F. Reich, 1997. An assessment of soil resources of Africa in relation to soil productivity. *Geoderma*, Vol. 77,1-18.

Eze C.C, 2011. Geo electrical studies for estimating aquifer hydraulic properties in Enugu State, Nigeria. *International Journal of the Physical Sciences*, Vol.6, No.14, 3319-3329.

Federal Department of Agricultural Land Resources (FEDLAR) 1990. The reconnaissance Soil Survey of Nigeria.

Gee, G.W., and D. Or, 2002. Particle Size analysis. p. 225-293 In D.J., and G.C Topps (eds.) *Methods of soil analysis. Dane Part 4. Physical methods. Soil Science Society of America Book Series No. 5*, ASA and SSSA. Madison, W.I.

Giesler, R., T. Andersson, L. Lovgren, and P. Persson, 2005. Phosphate sorption in aluminium- and iron-rich humus soil. *Soil Sci. Soc. Am.J.*, Vol., 69, 77-86.

Green V. T., M.A. Cavigelli, T.H. Dao and D.C. Flanagan, 2005. Soil physical properties and aggregate- associated C, N, and P distributions in organic and conventional cropping systems. *Soil Science*, Vol.170 No. 10,, 822- 831.

Grossman, R.B. and T.B. Reinsch, 2002. Bulk density and linear extensibility.p. 201-228. In M.J.H, Dane and G.C. Topp, (eds.) *Methods of the Soil Analysis part 4. Physical methods. Soil Sci. Soc. Am. Book Series No.5*, ASA and SSSA, Madison, WI.

Guan S., An N. Lui, J. Zong, N. He, Y. Shi, P. J. Zhang and N. He, 2018. Warming impacts on carbon, nitrogen and phosphorus distribution in soil water-stable aggregates. *Plant Soil Environment*, Vol. 64, No. 2, 64–69, URL: [http:// doi: 10.17221/715/2017-PSE](http://doi:10.17221/715/2017-PSE)

Haynes, R.J, and R.S., Swift, 1990. Stability of soils aggregates in relation to organic constituents and soil water content. *Journal of Soil Science*, Vol. 44,73-83.

Hendershort, W.H., H., Lalande, and M., Duquette, 1993. Soil reaction and exchangeable Acidity.p. 141-145. In M.R, Carter (ed.) *Soil sampling and methods of soil analysis*, Canadian Soc. Soil Sci. Lewis Publishers, London.

Igwe C. A, F.O.R., Akamigbo J.S.C and Mbagwu, 1995. Physical properties of soils of southeastern Nigeria and the role of some aggregating agents in their stability; *Soil Sci. Vol.160*, 431– 441.

Igwe C.A, 2000. Nutrient losses in runoff and eroded sediments from soils of central Eastern Nigeria. *Polish Journal of Soil Science*, Vol. 33 No.1, 67-75.

Igwe C.A, 2001. Water stable aggregates of Niger floodplain soils and their organic carbon Nitrogen and phosphorus distribution. *Agro Science. Journal of tropical agriculture, food, environment and extension*, Vol. 2. No.2, 52- 61.

Igwe C A., 2003. Erodibility of soils of the upper rainforest zone, southeastern Nigeria; *Land Degrad. Dev.* 14323–334.

Igwe, C. A., and K., Stahr, 2004. Waterstable aggregates of flooded Inceptisols from Southeastern Nigeria in relation to mineralogy and Chemical properties. *Australian Journal of Soil Research*, Vol.42,171-179.

Ihem, E.E., U.N., Nkwopara, B.N., Ndukwu, and S.U., Onwudike, 2017. Macro-Aggregate Stability of Soils under Selected Parent Materials in Imo State, Nigeria. *International Journal of Research in Agriculture and Forestry*, Vol. 4, No. 12, 24-32.

Iraj E., S. Reissand H. Rudolf Bork, 2009. A study of the relationship between land management and soil aggregate stability (case study near Albersdorf, Northern-Germany). *ARPJ Journal of Agricultural and Biological Science*, Vol. 4, 48-53.

Jiang X., Y. Hu, J.H. Bedel, D. Xie and A.L. Wright, 2011. Soil organic carbon and nutrient content in aggregate-size fractions of a subtropical rice soil under variable tillage. *Soil Use and Management*, Vol, 27, 28-35.

Kazman, Z., I. Sheinberg, M. Gal, 1983. Effects of low levels of exchangeable sodium and applied phosphor gypsum on the infiltration rate of various soils. *Soil Science*, Vol.135, 184-192.

Kemper, W.D., and R.C., Rosenau, 1986. Aggregate stability and size distribution p. 425-442. In A., Klute, (ed.) *Methods of soil analysis part 1, physical and mineralogical method*, 2nd Edition. American Society of Agronomy, Madison., W.I.

Lal R., 2009. Soil degradation as a reason for inadequate human nutrition. *Food Security*.1,45–57.

Landon, J.R., 1991. *Booker tropical soil manual: A hand book for soil Surveyors and agricultural land evaluation in the tropics*. Paperback edition, Long man Scientific and Technical.U.K.

Le Bissonnais., 1996. Aggregate stability and assessment of soil crustability and erodibility Theory and methodology. *European Journal of Soil Science*, Vol. .47, 425-437.

Le Bissonnais.Y., and M.J., Singer 1993. Seal formation, runoff and interrill erosion fromseventeen California soils. *Soil Sci. Soc of Amer. J.*, Vol. 41,954-960.

Mbagwu, J.S.C. and M.E. Obi, 2003. Land degradation, agricultural productivity and rural poverty. Environmental implications. Proceedings of the 28 the annual conference of Soil Science society of Nigeria. National Root Crop Research Institute, Umudike Umuahia, Nigeria Vol. 407: 1-11.

Mbagwu, J.S.C., A. Piccolo and M.O. Mbila, 1993. Water stability of aggregates of some tropical soils treated with humic substances. *Pedologie, Gent* Vol.43 No. 2, 269-284.

Nelson, D. N., and S.L. Sommers 1982. Total Carbon, organic carbon and matter, p, 539 - 579. In A.L., Page, R.H., Millerand D.K., Keeney (eds.) *Methods of soil analysis part 2* Amer. Soc. Agron. D.M.Madison, W. I.

Nigeria Meteorological Agency (NIMET) 2012. Seasonal Rainfall predictions and a Socio-Economic implications for Nigeria,1-22.

Nimmo, J.R, 1997. Modeling structural influences on soil water retention. *Soil Sci. Soc. Am. J.*, Vol. 61, 712-719.

Nimmo J.R., and K.S., Perkins, 2002. Aggregate stability and size distribution, p, 317-328. In J.H., Dane, and G.C., Topp (eds.) Methods of soil analysis, Part 4- physical methods: Madison, Wisconsin. Soil Science Society of America.

Obi M. 1982. Runoff and soil loss from an Oxisol in southeastern Nigeria under various management practices. *Agricultural Water Management*, Vol. 5, 193-203.

Ojanuga, A.G and A.I., Awojuola 1981. Characterisation and classification of the soils of the Jos plateau, Nigeria. *Nig J. Soil Sci.*, Vol., 10, 101 – 119.

Onweremadu, E.U., 2007. Availability of selected soil nutrients in relation to land use and landscape position. *Int. J. Soil Sci.*, Vol. 2, 128-134.

Onwudike, S. U., Onweremadu, E. U., Ihem, E. E., Agim, L.C., Osisi, A. F., Osuaku, S. K., and Azuh, P. O, 2016. Evaluation of Micronutrient Status of Soils under Three Land Use Types in Oyiabo, River State, Nigeria, *Futo Journal Series*. Vol. 2 No. 1, 32 - 40 .. URL: www.futojnls.org.

Opara-Nadi, O. A., 1988. Liming and organic matter interaction in two Nigerian Ultisols. Effect on soil pH, organic carbon and early growth of maize (*Zea mays*. L). Proceedings of the 16th annual conference of Soil Science Society of Nigeria. Minna, Niger State. Nov. 27 -30th 177- 198.

Razafimbelo, T.M., Albrecht, R., Oliver, T., Chevallier, L., Chapuis- Lardy, and C., Feller .2008. Aggregate associated-C and physical protection in a tropical clayey soil under Malagasy conventional and no-tillage systems. *Soil & Tillage Research*, Vol. 98, 140–149.

Sakin, E., 2012. Relationships between of Carbon, Nitrogen Stocks and Texture of the Harran Plain Soils in Southeastern Turkey. *Agricultural Academy. Bulgarian Journal of Agricultural Science*, Vol. 18 No. 4, 626- 634.

Soil Survey Staff, 2006. Soil quality information sheet; soil quality indicators aggregate stability. National Soil Survey Centre in collaboration with NRCS, USDA and the National Soil Tilth Laboratory. ARS, USDA.

Singer J.M., and W.D., Munns, 1996. *Soils .An Introduction*, 3rd Edition Prentice Hall-Inc. New Jersey.

Quirk, J.P., and J., Murray, 1991. Some physico-chemical aspects of soil structural stability. A review, In: *Modification of soil structure*, (Emerson W.W., R.D., Bonds Ed.), John Willy and Sons New York. 3-16.

Tisdall, J.M, and J.M., Oades, 1982. Organic matter and water stable aggregates in soils. *Journal of Soil Sci.*, Vol. .33, 141-163.

Thomas 1982. *Methods of soil analysis, part 2: Chemical and Microbiological properties*, 2nd Edition; Amer. Soc. Agro. Inc. Madison, Wisconsin, USA.

Reyment R.A, 1965. *Aspects of geology of Nigeria*. Ibadan University Press, Ibadan, 145.

Unamba-Opara I., 1982. Comparison of the carbon and Nitrogen contents and their relationships to other soil properties in some important soils of Southeastern Nigeria. *Beitrag trop. Landwirtsch. Veterinarmed*. Vol. 2, 167-176.

USDA, 1996. *Soil Quality Resource Concerns: Soil Erosion*. Soil Quality information sheet. USDA Natural resources Conservation service.

Uzoho, B. U. and N.N., Oti, 2005. Phosphorus adsorption characteristics of selected

Southeastern Nigerian Soils Journal of Agriculture, Food, Environment and Extension, Vol.4 1 , 50-55.

Verma, V. K, R. K. Setia, P.L. Sharma, S. Charanjit, and A. Kumar, 2005. Pedospheric variations in distribution of DPTA - extractable micronutrients in soils developed on different physiographic units in central parts of Punjab, India. Int. J. Agric. Biol., Vol. 7 No., 2, 243 – 246.

Wright A. L., and P. W. Inglett, 2009. Soil organic carbon and nitrogen and distribution of carbon-13 and Nitrogen-15 in aggregates of Everglades Histosols, Soil Science Society of American Journal, Vol., 73, 427-433, [URL:http://doi:10.2136/sssaj2008.0078](http://doi:10.2136/sssaj2008.0078).